


EMS System for Metropolitan Oklahoma City and Tulsa 2017 Medical Control Board Treatment Protocols



Approved 11/9/16, Effective 2/1/17, replaces all prior versions



FOR OPIATE USE, ADULT MUST HAVE SYS BP ≥ 100 mmHg; PEDIATRIC MUST HAVE SYS BP ≥ (70 + 2x age in years) mmHg

ADULT: FENTANYL 1 mcg/kg SLOW IVP/IM/IN, MAXIMUM DOSE 100 mcg. MAY REPEAT EVERY 10 MINUTES TO

MAXIMUM CUMULATIVE DOSE OF 3 mcg/kg or 250 mcg WHICHEVER IS LESSER.

OR

ADULT: MORPHINE SULFATE 2 - 4 mg SLOW IVP, MAY REPEAT 2 - 4 mg EVERY 5 MINUTES TO A TOTAL OF 10 mg. OR

ADULT: HYDROMORPHONE 0.5 - 1 mg SLOW IVP
MAY REPEAT EVERY 10 MINUTES TO MAXIMUM CUMULATIVE DOSE OF 2 mg.

PEDIATRIC: OLMCP ORDER ONLY

OLMCP CONSULT IF FURTHER ANALGESIA REQUIRED

CONTINUOUS ASSESSMENT & TREATMENT PER APPLICABLE PROTOCOL(S)