


EMS System for Metropolitan Oklahoma City and Tulsa 2017 Medical Control Board Treatment Protocols



EMS Section

Approved 11/9/16, Effective 2/1/17, replaces all prior versions



ADULT: IF CHEST PAIN IMPROVED WITH INITIAL NTG:

IF PT STILL HAVING CHEST PAIN AFTER 3 NTG ADMINISTRATIONS & IF SYS BP > 100 mmHg: ADDITIONAL NITROGLYCERIN PER PROTOCOL 16HH AND

MORPHINE SULFATE 2 mg SLOW IVP, MAY REPEAT EVERY 5 MIN TO A TOTAL OF 10 mg. OR

FENTANYL 0.5 mcg/kg SLOW IVP/IM/IN, MAXIMUM DOSE 50 mcg. MAY REPEAT EVERY 10 MINUTES TO MAXIMUM CUMULATIVE DOSE OF 1.5 mcg/kg or 125 mcg WHICHEVER IS LESSER. ŌR

> HYDROMORPHONE 0.25 mg SLOW IVP MAY REPEAT EVERY 10 MINUTES TO MAXIMUM CUMULATIVE DOSE OF 1 mg.

ADULT: IF CHEST PAIN NOT IMPROVED WITH INITIAL NTG: MAY TREAT WITH OPIATE OPTION ABOVE IF INDICATED PEDIATRIC: CONSULT FOR TREATMENT DIRECTIVE(S) IF INDICATED