



STATE OF OKLAHOMA 2013 EMERGENCY MEDICAL SERVICES PROTOCOLS



TREATMENT PRIORITIES

- Vital signs
- IV Access
- Analyze rhythm
- Rhythm specific treatment
- Consult OLMC for pediatric pts
- If protocol measures DO NOT control rate of adult PSVT, or convert WIDE COMPLEX TACHYCARDIA, **CONTACT OLMC**
- OLMC Adult options include:
 - Narrow Complex Tachycardia**
 - DILTIAZEM 0.25 mg/kg SLOW IVP over 2 mins
 - AMIODARONE 150mg SLOW IVP/IVPB over 10 mins
 - SYNCHRONIZED** CARDIOVERSION
 - MONITOR/TRANSPORT
 - Wide Complex Tachycardia**
 - Repeat AMIODARONE 150mg SLOW IVP/IVPB over 10 mins
 - SYNCHRONIZED** CARDIOVERSION
 - ADENOSINE 12 mg RAPID IVP IF REGULAR & MONOMORPHIC
 - LIDOCAINE up to 1 mg/kg IVP at < 50 mg/min
 - MONITOR/TRANSPORT

UNSTABLE (SYMPTOMATIC) tachycardia is defined by any of the following symptoms:

- Dyspnea
- Chest pain
- Weakness
- Altered mental status
- Hypoxemia
- Pulmonary edema

